



WHY IS MY LEVOTHYROXINE PACKAGED IN A BLISTER PACK?



EUTHYROX® (levothyroxine sodium) tablets are available in the following strengths: 25 mcg, 50 mcg, 75 mcg, 88 mcg, 100 mcg, 112 mcg, 125 mcg, 137 mcg, 150 mcg, 175 mcg and 200 mcg. (Illustration above is for the 100 mcg strength only.) Each box and inner blister are color coded to match the corresponding levothyroxine strength.

We are excited to offer you the world's leading brand¹ of levothyroxine (EUTHYROX®) as the substitute for levothyroxine and Synthroid® prescriptions. Some of the benefits of this include the following:

- EUTHYROX® (levothyroxine sodium) blister packaging **protects** the tablets from light and moisture (humidity), either of which can cause any levothyroxine tablet to degrade and lose some of its potency².
- With EUTHYROX® (levothyroxine sodium) tablets, you can rest assured that the strength of the first tablet consumed is the same as the last³.
- With EUTHYROX® (levothyroxine sodium) tablets, you can easily see exactly how many tablets are left before you need to obtain a refill.
- Taking your prescription as directed by your healthcare provider every day and having it refilled on-time may result in more consistent thyroid test results⁴.
- The outer packaging of EUTHYROX® has braille imprinting for the visually impaired.

Additional product features:

- All strengths of EUTHYROX® (levothyroxine sodium) tablets (25 mcg, 50 mcg, 75 mcg, 88 mcg, 100 mcg, 112 mcg, 125 mcg, 137 mcg, 150 mcg, 175 mcg and 200 mcg) contain no dyes, lactose or gluten, which may reduce the potential for adverse effects in patients with sensitivities to those ingredients.
- Each tablet is embossed with the actual dosage strength (see below).

25 mcg	50 mcg	75 mcg	88 mcg	100 mcg	112 mcg	125 mcg	137 mcg	150 mcg	175 mcg	200 mcg

Refer to the IMPORTANT SAFETY INFORMATION located on the reverse side of this page. If you have any questions about your EUTHYROX® prescription, talk to your pharmacist.

¹ Data on file – Merck KGaA Darmstadt, Germany – IQVIA Midas Health Q3 2018

² Benvenga et al. (2017). Refractory Hypothyroidism Due to Improved Storage of Levothyroxine Tablets. *Frontiers in Endocrinology, Vol 8, Article 155*

³ Data on file – Provell Pharmaceuticals, LLC

⁴ Shrestha et al. (2018) Medication Adherence to Levothyroxine Therapy among Hypothyroid Patients and their Clinical Outcomes with Special Reference to Thyroid Function Parameters. *Kathmandu Univ Med J (KUMJ)* Apr-Jun; 16(62) 129-137 <https://www.ncbi.nlm.nih.gov/pubmed/30636753>

All other trademarks, registered or unregistered, are the property of their respective owners.

DESCRIPTION OF USE:

EUTHYROX® (levothyroxine sodium) tablets, is a prescription, thyroid hormone that is used to treat a condition called hypothyroidism. It is usually taken by mouth and should be taken as directed by your healthcare provider. It is meant to replace a hormone that is normally made by your thyroid gland. Generally, thyroid replacement treatment is to be taken for life. EUTHYROX should not be used to treat noncancerous growths or enlargement of the thyroid in patients with normal iodine levels, or in cases of temporary hypothyroidism caused by inflammation of the thyroid gland (thyroiditis).

IMPORTANT SAFETY INFORMATION

- **Thyroid hormones, including EUTHYROX, should not be used either alone or in combination with other drugs for the treatment of obesity or weight loss. Larger doses may result in serious or even life-threatening events, especially when used in combination with certain other drugs used to reduce appetite.**
- Do not use EUTHYROX if you have uncorrected adrenal problems.
- Use EUTHYROX only as directed by your prescriber. Take EUTHYROX as a single dose, preferably on an empty stomach, one-half to one hour before breakfast.
- Tell your doctor if you are pregnant or breastfeeding or are thinking of becoming pregnant while taking EUTHYROX. Your dose of EUTHYROX may need to be increased during your pregnancy.
- Taking too much levothyroxine has been associated with increased bone loss, especially in women after menopause.
- Once your prescriber has found your specific EUTHYROX dose, it is important to have lab tests completed, as ordered by your doctor, to ensure that you are on the correct dose.
- Talk with your healthcare provider about your diet as foods like soybean flour, cottonseed meal, walnuts, and dietary fiber may cause your body to absorb less EUTHYROX from the gastrointestinal tract. Grapefruit juice may cause your body to absorb less levothyroxine and may reduce its effect. If you eat these types of food, your dose of EUTHYROX may need to be adjusted.
- Products such as iron and calcium supplements and antacids can lower your body's ability to absorb levothyroxine, so EUTHYROX should be taken 4 hours before or after taking these products.
- It may take several weeks before you notice an improvement in your symptoms.
- Tell your doctor if you are taking any other drugs, including prescription and over-the-counter products.
- Tell your doctor about any other medical conditions you may have, especially heart disease, diabetes, blood clotting problems, and adrenal or pituitary gland problems. The dose of other drugs you may be taking to control these conditions may have to be changed while you are taking EUTHYROX. If you have diabetes, check your blood sugar levels and/or the glucose in your urine, as ordered by your doctor, and immediately tell your doctor if there are any changes. If you are taking blood thinners, your blood clotting status should be checked frequently.
- Tell your doctor or dentist that you are taking EUTHYROX before any surgery.
- Tell your doctor if you develop any of the following symptoms: rapid or abnormal heartbeat, chest pain, shortness of breath, leg cramps, headache, nervousness, irritability, sleeplessness, shaking, change in appetite, weight gain or loss, vomiting, diarrhea, increased sweating, difficulty tolerating heat, fever, changes in menstrual periods, swollen red bumps on the skin (hives) or skin rash, or any other unusual medical event.
- Partial hair loss may occur during the first few months you are taking EUTHYROX.

This is the most important safety information you should know about EUTHYROX. For more information, talk with your doctor or healthcare provider.

Please refer to the full prescribing information located within your EUTHYROX prescription box or visit <https://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?setid=bc3c6830-ba68-4ba3-9896-a12d0ad0b275>

You are encouraged to report negative side effects of prescription drugs to the FDA. You can do this by contacting Provell Pharmaceuticals, LLC at 1-888-899-7041 or the FDA at [1-800-FDA-1088](https://www.fda.gov/medwatch) or you can visit www.fda.gov/medwatch.

Provell Pharmaceuticals, LLC - PP-RG-00003



NOW AVAILABLE